

Multiple Choice Questions

1. Agomelatine acts on the following receptors:

MT1	T	F
MT2	T	F
MT3	T	F
5-HT _{2a}	T	F
5-HT _{2c}	T	F

2. The following agents are recognised treatment options for rapid cycling BPAD:

Risperidone	T	F
Quetiapine	T	F
Carbamazepine	T	F
Lamotrigine	T	F
Zonisamide	T	F

3. St. John's wort (*Hypericum perforatum*) is associated with:

Serotonin re-uptake inhibition	T	F
Noradrenaline re-uptake inhibition	T	F
Dopamine re-uptake inhibition	T	F
Inducing hypomanic episodes	T	F
Increasing plasma concentrations of warfarin	T	F

4. In psychotic depression:

The most effective treatment option is MAOIs	T	F
Mifepristone improves psychotic more than depressive symptoms	T	F
Treatment response is approximately equal to non-psychotic depression	T	F
Is best treated with an antidepressant and antipsychotic agent	T	F
rTMS has proven efficacy	T	F

5. Vagal Nerve Stimulation in depression is associated with

An approximate 1% risk of vocal palsies	T	F
An approximate 1% risk of voice alteration	T	F
Voice alteration seldom persists with continued treatment	T	F
The induction of rapid-cycling BPAD in some individuals	T	F
A stronger evidence base than for the treatment of BPAD	T	F

Answers: